

From ordered semigroups to ordered hypersemigroups

Niovi KEHAYOPULU*

Department of Mathematics, University of Athens, 15784 Panepistimiopolis, Greece

Abstract: We wrote this paper as an example to show the way we pass from ordered semigroups to ordered hypersemigroups.

Key words: ordered hypersemigroup, right (left) ideal, bi-ideal, quasi-ideal, regular, intra-regular

1 Introduction and prerequisites

For an ordered semigroup $(S, \cdot, \leq)$ and a subset A of S , we denote by $(A]$ the subset of S defined by $(A] := \{t \in S \mid t \leq a \text{ for some } a \in A\}$. If $(S, \cdot, \leq)$ is an ordered semigroup, a nonempty subset A of S is called a right (resp. left) ideal of S if (1) $AS \subseteq A$ (resp. $SA \subseteq A$) and (2) if $a \in A$ and $S \ni b \leq a$, then $b \in A$, that is, if $(A] = A$ (cf. e.g. [3]). An ordered semigroup $(S, \cdot, \leq)$ is called regular [2] if for every $a \in S$ there exists $x \in S$ such that $a \leq axa$. It is called intra-regular [3] if for every $a \in S$ there exist $x, y \in S$ such that $a \leq xa^2y$. A subset B of an ordered semigroup $(S, \cdot, \leq)$ is called a bi-ideal of S [1] if (1) $BSB \subseteq B$ and (2) if $a \in B$ and $S \ni b \leq a$, then $b \in B$. A subset Q of S is called a quasi-ideal of S [4] if (1) $(QS) \cap (SQ) \subseteq Q$ and (2) if $a \in Q$ and $S \ni b \leq a$, then $b \in Q$. We have seen in [6], that an ordered semigroup S is regular if and only if for every right ideal A and every left ideal B of S , we have $A \cap B = (AB]$, equivalently, $A \cap B \subseteq (AB]$. In [8] we have shown that an ordered semigroup S is intra-regular if and only if for every right ideal A and every left ideal B of S we have $A \cap B \subseteq (BA]$. In [5], we proved that if S is a regular ordered semigroup, then B is a bi-ideal of S if and only if there exist a right ideal R and a left ideal L of S such that $B = (RL]$. We have seen in [6] that an ordered semigroup S is

*Correspondence: nkehayop@math.uoa.gr

2010 *AMS Mathematics Subject Classification*: Primary 06F99, Secondary 06F05

regular if and only if the ideals of S are idempotent and for every right ideal A and every left ideal B of S , the set $(AB]$ is a quasi-ideal of S . In [7] we studied some of the above results for hypersemigroups. In the present paper we examine the results on ordered semigroups mentioned above for ordered hypersemigroups. The results of ordered semigroups can be transferred to ordered hypersemigroups in the way indicated in the present paper. For the definitions and notations not given in the present paper we refer to [7]. The concept of an ordered semigroup can be transferred in a natural way to ordered hypersemigroups, for this concept we will refer to [9], and we will give further information about it in a next paper. In the proofs we tried to use sets instead of elements to show their pointless character.

2 Main results

Based on ordered semigroups, we first introduce the concepts of regular and intra-regular hypersemigroups and we characterize them in terms of right and left ideals. The concept of regularity of ordered semigroups is naturally transferred to an hypersemigroup H in the following definition

Definition 1. An ordered hypersemigroup $(H, \circ, \leq)$ is called *regular* if for every $a \in H$ there exist $x, y \in H$ such that $y \in (a \circ x) * \{a\}$ and $a \leq y$.

Proposition 2. Let H be an ordered hypersemigroup. The following are equivalent:

- (1) H is regular.
- (2) $a \in (\{a\} * H * \{a\})$ for every $a \in H$.
- (3) $A \subseteq (A * H * A)$ for every $A \in \mathcal{P}^*(H)$.

Definition 3. Let H be an ordered hypergroupoid. A nonempty subset A of H is called a *right* (resp. *left*) *ideal* of H if

- (1) $A * H \subseteq A$ (resp. $H * A \subseteq A$) and
- (2) if $a \in A$ and $H \ni b \leq a$, then $b \in A$, that is, if $(A] = A$.

Lemma 4. Let H be an ordered hypergroupoid and $A, B \in \mathcal{P}^*(H)$. Then we have

$$(A] * (B] \subseteq (A * B).$$

Lemma 5. For an hypersemigroup H and a nonempty subset A of H , we denote by $R(A)$ (resp. $L(A)$), the right (resp. left) ideal of H generated by A , and by $I(A)$ the ideal of H generated by A . We have

$$\begin{aligned} R(A) &= \left(A \cup (A * H) \right], \quad L(A) = \left(A \cup (H * A) \right], \\ I(A) &= \left(A \cup (H * A) \cup (A * H) \cup (H * A * H) \right]. \end{aligned}$$

Lemma 6. Let H be an hypergroupoid. If A is a right ideal and B a left ideal of H , then $A \cap B \neq \emptyset$.

Lemma 7. Let H be an ordered hypergroupoid and $A, B \in \mathcal{P}^*(H)$. Then we have

$$(A * B] = \left((A] * (B] \right] = \left((A] * B \right] = \left(A * (B] \right].$$

Theorem 8. An ordered hypersemigroup H is regular if and only if for every right ideal A and every left ideal B of H , we have

$$A \cap B = (A * B], \text{ equivalently, } A \cap B \subseteq (A * B].$$

Proof. Let H be regular, A a right ideal and B a left ideal of H . Since $A \cap B \neq \emptyset$ and H is regular, by Proposition 2, we have

$$\begin{aligned} A \cap B &\subseteq \left((A \cap B) * H * (A \cap B) \right] \subseteq \left((A * H) * B \right] \subseteq (A * B] \\ &\subseteq (A * H] \cap (H * B] \subseteq (A] \cap (B] = A \cap B. \end{aligned}$$

Thus we have $A \cap B = (A * B]$.

Suppose now that $A \cap B \subseteq (A * B]$ for every right ideal A and every left ideal B of H , and let $A \in \mathcal{P}^*(H)$. Then we have

$$\begin{aligned} A &\subseteq R(A) \cap L(A) \subseteq \left(R(A) * L(A) \right] = \left(\left(A \cup (A * H) \right] * \left(A \cup (H * A) \right] \right] \\ &\subseteq \left(\left(A \cup (A * H) \right) * \left(A \cup (H * A) \right) \right] \text{ (by Lemma 7)} \\ &= \left((A * A) \cup (A * H * A) \cup (A * H * H * A) \right] \\ &= \left((A * A) \cup (A * H * A) \right]. \end{aligned}$$

Then

$$\begin{aligned} A * A &\subseteq \left((A * A) \cup (A * H * A) \right] * (A] \subseteq \left((A * A * A) \cup (A * H * A * A) \right] \\ &\subseteq (A * H * A]. \end{aligned}$$

Then we have

$$A \subseteq \left((A * H * A] \cup (A * H * A) \right] = \left((A * H * A] \right] = (A * H * A]$$

and, by Proposition 2, H is regular. $\square$

The concept of intra-regularity of ordered semigroups is naturally transferred to an hypersemigroup H in the following definition.

Definition 9. An ordered hypersemigroup H is called *intra-regular* if for every $a \in H$ there exist $x, y, t \in H$ such that

$$t \in (x \circ a) * (a \circ y) \text{ and } a \leq t.$$

Proposition 10. Let H be an ordered hypersemigroup. The following are equivalent:

- (1) H is intra-regular.
- (2) $a \in (H * \{a\} * \{a\} * H]$ for every $a \in H$.
- (3) $A \subseteq (H * A * A * H]$ for every $A \in \mathcal{P}^*(H)$.

Theorem 11. An ordered hypersemigroup H is intra-regular if and only if for every right ideal A and every left ideal B of H , we have

$$A \cap B \subseteq (B * A].$$

Proof. $\implies$. Let A be a right ideal and B a left ideal of H . By Lemma 6, $A \cap B \in \mathcal{P}^*(H)$. Since H is intra-regular, by Proposition 10, we have

$$\begin{aligned} A \cap B &\subseteq \left(H * (A \cap B) * (A \cap B) * H \right] \\ &\subseteq \left((H * B) * (A * H) \right] \\ &\subseteq B * A. \end{aligned}$$

$\impliedby$. Just for a change, let us prove it using elements. The same can be proved using sets, as we did in Theorem 8. Let $a \in H$. By hypothesis, we have

$$\begin{aligned} a &\in R(a) \cap L(a) \subseteq \left(L(a) * R(a) \right] \\ &= \left(\left(\{a\} \cup (H * \{a\}) \right) * \left(\{a\} \cup (\{a\} * H) \right) \right] \\ &\subseteq \left(\left(\{a\} \cup (H * \{a\}) \right) * \left(\{a\} \cup (\{a\} * H) \right) \right] \text{ (by Lemma 7)} \\ &= \left((a \circ a) \cup (H * \{a\} * \{a\}) \cup (\{a\} * \{a\} * H) \cup (H * \{a\} * \{a\} * H) \right]. \end{aligned}$$

Then $a \in (a \circ a]$ or $a \in (H * \{a\} * \{a\})$ or $a \in (\{a\} * \{a\} * H]$ or $a \in (H * \{a\} * \{a\} * H]$.

If $a \in (a \circ a]$, then

$$\begin{aligned} a &\in \left((a \circ a) * (a \circ a) \right] \subseteq \left((a \circ a) * (a \circ a) \right] \text{ (by Lemma 7)} \\ &\subseteq (H * \{a\} * \{a\} * H] \end{aligned}$$

then, by Proposition 10, H is intra-regular. If $a \in (H * \{a\} * \{a\})$, then

$$\begin{aligned} a &\in \left(H * (H * \{a\} * \{a\}) * \{a\} \right] = (H * H * \{a\} * \{a\} * \{a\}) \\ &\subseteq (H * \{a\} * \{a\} * H]. \end{aligned}$$

If $a \in (H * \{a\} * \{a\})$ or $a \in (H * \{a\} * \{a\} * H]$ in a similar way, we have $a \in (H * \{a\} * \{a\} * H]$. In any case, H is intra-regular. $\square$

Below we first transfer the concept of bi-ideals from ordered semigroups to ordered hypersemigroups, then we characterize the bi-ideals of ordered hypersemigroups in terms of right and left ideals.

Definition 12. Let $(H, \circ, \leq)$ be an ordered hypersemigroup. An element $B \in \mathcal{P}^*(H)$ is called a *bi-ideal* of H if

- (1) $B * H * B \subseteq B$ and
- (2) if $a \in B$ and $H \ni b \leq a$, then $b \in B$, that is if $(B] = B$.

Proposition 13. Let H be an ordered hypersemigroup. If C is a right ideal and $D \in \mathcal{P}^*(H)$, then the set $B = (C * D]$ is a bi-ideal of H .

Proof. We have

$$\begin{aligned} B * H * B &= (C * D] * (H] * (C * D] \\ &\subseteq \left((C * D) * H \right] * (C * D] \text{ (by Lemma 4)} \\ &\subseteq \left((C * D) * H * (C * D) \right] \text{ (by Lemma 7)} \\ &= \left((C * (D * H * C) * D) \right] \subseteq \left((C * H) * D \right] \\ &\subseteq (C * D] = B. \end{aligned}$$

$\square$

In a similar way we have the following proposition

Proposition 14. Let H be an ordered hypersemigroup. If C is a left ideal and $D \in \mathcal{P}^*(H)$, then the set $B = (C * D]$ is a bi-ideal of H .

By Propositions 13 and 14 we have the following

Proposition 15. *Let H be an ordered hypersemigroup. If C is a right ideal and D a left ideal of H , then the set $B = (C * D]$ is a bi-ideal of H .*

Proposition 16. *Let H be a regular ordered hypersemigroup and B a bi-ideal of H . Then there exist a right ideal C and a left ideal D of H such that $B = (C * D]$.*

Proof. Since B is a bi-ideal of H , we have $B * H * B \subseteq B$. Since H is regular, by Proposition 2, we have $B \subseteq (B * H * B]$. Thus we have $B = (B * H * B]$.

On the other hand,

$$\begin{aligned} R(B) * L(B) &= \left((B \cup (B * H)) \right] * \left((B \cup (H * B)) \right] \\ &\subseteq \left(\left((B \cup (B * H)) \right) * \left(B \cup (H * B) \right) \right] \quad (\text{by Lemma 7}) \\ &= \left((B * B) \cup (B * H * B) \cup (B * H * H * B) \right] \\ &= \left((B * B) \cup (B * H * B) \right]. \end{aligned}$$

Since

$$B * B = (B] * (B * H * B] \subseteq (B * B * H * B] \subseteq (B * H * B],$$

we have

$$\begin{aligned} R(B) * L(B) &\subseteq \left((B * H * B] \cup (B * H * B) \right] = \left((B * H * B] \right] \\ &= (B * H * B] = B, \end{aligned}$$

so $\left(R(B) * L(B) \right] \subseteq (B] = B$. On the other hand, we have

$$\begin{aligned} B &= (B * H * B] \subseteq \left(\left(R(B) * H \right) * L(B) \right] \\ &\subseteq \left(R(B) * L(B) \right] \quad (\text{since } R(B) \text{ is a right ideal of } H). \end{aligned}$$

Hence we obtain $B = \left(R(B) * L(B) \right]$, where $R(B)$ is a right ideal and $L(B)$ is a left ideal of H . $\square$

By Propositions 15 and 16 we have the following theorem

Theorem 17. *Let H be a regular ordered hypersemigroup. An element $B \in \mathcal{P}^*(H)$ is a bi-ideal of H if and only if there exist a right ideal C and a left ideal D of H such that $B = (C * D]$.*

In the rest of the paper we characterize the regular hypersemigroups in terms of right ideals, left ideals and quasi-ideals. The concept of quasi-ideals of ordered semigroups is naturally transferred to ordered hypersemigroups as follows:

Definition 18. Let $(H, \circ, \leq)$ be an ordered hypersemigroup. An element $Q \in \mathcal{P}^*(H)$ is called a *quasi-ideal* of H if

- 1) $(Q * H] \cap (H * Q] \subseteq Q$ and
- 2) if $a \in Q$ and $H \ni b \leq a$, then $b \in Q$.

We first give the following definition, proposition.

Definition 19. Let H be an ordered hypersemigroup. An element $A \in \mathcal{P}^*(H)$ is called *subidempotent* if $(A * A] \subseteq A$. It is called *idempotent* if $(A * A] = A$.

Proposition 20. *If H is an ordered hypergroupoid, then the right ideals and the left ideals of H are subidempotent. In particular, if H is a regular ordered hypersemigroup, then the right ideals and the left ideals of H are idempotent.*

Theorem 21. *An ordered hypersemigroup H is regular if and only if the right ideals and the left ideals of H are idempotent and for every right ideal A and every left ideal B of H , the set $(A * B]$ is a quasi-ideal of H .*

Proof. $\implies$. By Proposition 20, the right and the left ideals of H are idempotent. Let now A be a right ideal and B a left ideal of H . Since H is regular, by Theorem 8, we have $A \cap B = (A * B]$. It is enough to prove that $A \cap B$ is a quasi-ideal of H . First of all, by Lemma 6, $A \cap B \in \mathcal{P}^*(H)$. Moreover,

$$\begin{aligned} \left((A \cap B) * H \right] \cap \left(H * (A \cap B) \right] &\subseteq (A * H] \cap (H * B] \\ &\subseteq (A] \cap (B] = A \cap B, \end{aligned}$$

and if $x \in A \cap B$ and $H \ni y \leq x$ then, since $y \leq x \in A$ and A is an ideal of H we have $y \in A$ and since $y \leq x \in B$, we have $y \in B$, so $y \in A \cap B$. Thus $A \cap B$ is a quasi-ideal of H , and so is $(A * B]$.

$\impliedby$. Let $A \in \mathcal{P}^*(H)$. Since $R(A)$ is a right ideal of H , it is idempotent, and we have

$$\begin{aligned} A &\subseteq R(A) = \left(R(A) * R(A) \right] = \left(\left(A \cup (A * H) \right] * \left(A \cup (A * H) \right] \right] \\ &= \left(\left(A \cup (A * H) \right) * \left(A \cup (A * H) \right) \right] \text{ (by Lemma 7)} \end{aligned}$$

$$\begin{aligned}
&= \left((A * A) \cup (A * H * A) \cup (A * A * H) \cup (A * H * A * H) \right) \\
&\subseteq (A * H].
\end{aligned}$$

Since $L(A)$ is a left ideal of H , in a similar way, we have $A \subseteq (H * A]$. Thus we get $A \subseteq (A * H] \cap (H * A]$. Since $(A * H]$ is a right ideal and $(H * A]$ a left ideal of H , they are idempotent, and we have

$$\begin{aligned}
(A * H] \cap (H * A] &= \left((A * H] * (A * H] \right) \cap \left((H * A] * (H * A] \right) \\
&= \left((A * H) * (A * H) \right) \cap \left((H * A) * (H * A) \right) \quad (\text{by Lemma 7}) \\
&= \left((A * H * A) * H \right) \cap \left(H * (A * H * A) \right) \\
&= \left((A * H * A] * H \right) \cap \left(H * (A * H * A] \right) \quad (\text{by Lemma 7}).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
(A * H * A] &= \left(A * (H * H) * A \right) \quad (\text{since } H \text{ is a left ideal of } H) \\
&= \left(A * (H * H) * A \right) \quad (\text{by Lemma 7}) \\
&= \left((A * H) * (H * A) \right) \\
&= \left((A * H] * (H * A] \right) \quad (\text{by Lemma 7}).
\end{aligned}$$

Since $(A * H]$ is a right ideal and $(H * A]$ a left ideal of H , by hypothesis, the set $\left((A * H] * (H * A] \right)$ is a quasi-ideal of H , and so is $(A * H * A]$. Hence we obtain

$$\left((A * H * A] * H \right) \cap \left(H * (A * H * A] \right) \subseteq (A * H * A],$$

then $A \subseteq (A * H * A]$ and by Proposition 2, H is regular. $\square$

References

- [1] Kehayopulu N. On completely regular *poe*-semigroups. Math. Japon. 1992; 37: 123–130.
- [2] Kehayopulu N. On regular duo ordered semigroups. Math. Japon. 1992; 37: 535–540.
- [3] Kehayopulu N. On prime, weakly prime ideals in ordered semigroups. Semigroup Forum 1992; 44: 341–346.

- [4] Kehayopulu N. On regular, intra-regular ordered semigroups. Pure Math. Appl. (PU.M.A.) 1993; 4: 447–461.
- [5] Kehayopulu N. Note on bi-ideals in ordered semigroups. Pure Math. Appl. (PU.M.A.) 1995; 6: 333–344.
- [6] Kehayopulu N. On regular ordered semigroups. Math. Japon. 1997; 43: 549–553.
- [7] Kehayopulu N. On hypersemigroups. Pure Math. Appl. 2015; 25: 151–156.
- [8] Kehayopulu N, Lajos S, Tsingelis M. On regular, intra-regular ordered semigroups. Pure Math. Appl. 1993; 4: 447–461.
- [9] Pibaljommee B, Davvaz B. Characterizations of (fuzzy) bi-hyperideals in ordered semihypergroups. Journal of Intelligent & Fuzzy Systems 2015; 28: 2141–2148.